9000165

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Northrup King Co.

Chereas. There has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-CLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, R IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT TY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

SOYBEAN

'S28-18'

In Testimony Whereot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C. this 31st day of the year of our Lord one thousand nine hundred and ninety-one.

Hward Madigin

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, DIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

FORM APPROVED: OMB 0581-0055, Expires 1/31/91

U.S. DEPARTMENT OF A AGRICULTURAL MARKE APPLICATION FOR PLANT VARIET (Instructions on	TING SERVICE Y PROTECTION	I CERTIFICATE	deler certii Intor	tication is required in order to mine it a plant variety protection licate is to be issued (7 U.S.C. 2421). mation is held confidential until licate is issued (7 U.S.C. 2426).
NAME OF APPLICANT(S) (as it is to appear on the Certificate)		2. TEMPORARY DESIGNATION EXPERIMENTAL NO.	OR 3. V	ARIETY NAME
Northrup King Co.		X8928, J405545	2	328-18
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)		5 PHONE (Include area code)		FOR OFFICIAL USE ONLY
P. O. Box 959		r en	PVPO	NUMBER
Minneapolis, MN 55440		612-593-7333		9000165
			F	May 4, 1990
6. GENUS AND SPECIES NAME	7. FAMILY NAME (Botania	cal)	, N	Time
Glycine max	Leguminosae		G	A.M P.M.
8 CROP KIND NAME (Common Name)	9	DATE OF DETERMINATION	F E	Filing and Examination Fee:
Soybean		September, 1987	E S	Bala
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGA	NIZATION (Corporation, par		R	May 4,1990
Corporation			. C	Certificate Fee:
11. IF INCORPORATED, GIVE STATE OF INCORPORATION	12. DA	ATE OF INCORPORATION	Ţ	\$ 250.
Delaware		1976	E	Dec. 9 1991
Robert W. Romig Northrup King Co. P. O. Box 959 Minneapolis, MN 55440 14 CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Folia X) Exhibit A, Origin and Breeding History of the Variety b. X Exhibit B, Novelty Statement. c. X Exhibit C, Objective Description of Variety d. X Exhibit D, Additional Description of Variety. e. X Exhibit E, Statement of the Basis of Applicant's Ownersh t. X Seed Sample (2,500 viable untreated seeds). Date Seed g. X Filing and Examination Fee (\$2,150) made payable to " 15 DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SO Protection Act.) YES (If "YES." answer items 16 and 17 be NUMBER OF GENERATIONS? YES X NO 18 DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VARIETY YES." Through Plant Variety Protection Act	tip. I Sample mailed to Plant Vicessurer of the United Stock By VARIETY NAME ONLE WORLD IN TO 17. IF "YES" T	Variety Protection Office	D? (See secti	
X NO 19 HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR M				
YES (II "YES," give names of countries and dates)	BONNETED IN THE U.S. UNI	omen dodernied?		
20 The applicant(s) declare(s) that a viable sample of basic so request in accordance with such regulations as may be app. The undersigned applicant(s) is (are) the owner(s) of this uniform, and stable as required in section 41, and is entitle Applicant(s) is (are) informed that false representation her	licable. sexually reproduceded to protection under t	novel plant variety, and be he provisions of section 42 o	elieve(s) th f the Plant	nat the variety is distinct,
SIGNATURE OF APPLICANT (Owner(S))	CAPACITY OR	TITLE	T C	DATE
Aber M/ Normy	Vice f	PEGIDENI, RETERRE	cH	May 2, 1990
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR	TITLE		DATE

FORM CSSD-470 (5-89) Edition of FORM LS-470, 3-86, is obsolete

EXHIBIT A

Origin and Breeding History of the Variety

The soybean variety 'S28-18' is derived from the cross 'S30-31' \times 'A3127' which was made in 1981 by the soybean research group at St. Joseph, Illinois. The F1 and F2 generations were grown at the Northrup King Research Center near Waimea, Kauai, Hawaii in the winter of 1981-82. The F3 generation was grown in the field at St. Joseph in 1982, the F4 at Waimea in the winter of 1982-83. to F4 were all advanced by harvesting 2 to 4 pods from each plant. The F5 was planted at St. Joseph in the spring of 1983 and in the fall 100 random plants were harvested and threshed individually. The seed from these plants were grown in F6 progeny rows in 1984. One of these, numbered J405545, was selected based on agronomic appearance to be tested in a preliminary yield trial in 1985. has been tested at several midwest locations from 1986 to 1989 and found to yield well in comparison to other Maturity Group II cultivars. The line was tested from 1987 to 1989 for reaction to iron-deficiency chlorosis on calcareous soil near Harcourt in North Central Iowa and found to give an intermediate reaction. tested for reaction to Phytophthora megasperma and found to be susceptible. During the evaluation of J405545, the descriptive characteristics tawny pubescence, tan pods, and black hilum were identified and confirmed to be homogeneous, but flower color was found to be heterogeneous. In the winter of 1986-87 300 seeds were planted at Waimea and all purple flowered plants were removed. At harvest, 100 plants were threshed individually and their progeny planted at St. Joseph in 1987 to monitor variety variability and to produce Pedigree Seed. A few off-type plants, assumed to have come from outcrosses or mixtures, and two rows which marginally different maturity were removed. The remaining rows were uniform for white flower color and other characteristics and were bulked to produce Pedigree Seed. The resulting line was subsequently named S28-18. This seed was planted in 1988 to produce Breeder Seed. The increase block was rogued intensively at flowering and prior to harvest. A few plants with grey pubescence were removed. These were assumed to have resulted from admixture.

Foundation Seed of \$28-18 was grown in 1989. A few plants with grey pubescence were removed. The Iowa Crop Improvement Association inspected the field and found it to meet the standards for Foundation Seed. The National Soybean Variety Review Board approved the variety for Certification on December 7, 1989.

S28-18 is a stable and uniform variety except that it may contain up to 2% seed with hilum color other than black. In six years of testing and three cycles of seed increase, no variants other than those described previously have been observed.

Varietal purity will be maintained by use of progeny rows as needed.

EXHIBIT B

Novelty Statement for the Variety

Soybean variety \$28-18 is most similar to Preston. It can be differentiated from Preston based on flower color and pod color. \$28-18 has white flowers and tan pods while Preston has purple flowers and brown pods.

EXHIBIT C (Soybean)

U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
LIVESTOCK, MEAT, GRAIN & SEED DIVISION
PLANT VARIETY PROTECTION OFFICE
BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY

TEMPORARY DESIGNATION NAME S28-1	SOYB	EAN (Glycine max L.)		
ADDRESS (Street and No., or R.F.D. No., City, State, and Zip Code) P. O. Box 959 Minneapolis, MN 55440 Attention: R.W. Romig Choose the appropriate response which characterizes the variety in the features described below. When the number of significant digits in your answer is fewer than the number of boxes provided, place a zero in the first box when number is 9 or less (e.g., o 9). 1. SEED SHAPE: 2	NAME OF APPLICANT(S)	TEMPORARY DESIGNATION	VARIETY NAME	· · · · · · · · · · · · · · · · · · ·
Minneapolis, MN 55440 Attention: R.W. Romig Choose the appropriate response which characterizes the variety in the features described below. When the number of significant digits in your answer is fewer than the number of boxes provided, place a zero in the first box when number is 9 or less (e.g., 9). 1. SEED SHAPE:	Northrup King Co.		S28-18	
Choose the appropriate response which characterizes the variety in the features described below. When the number of significant digits in your answer is fewer than the number of boxes provided, place a zero in the first box when number is 9 or less (e.g., 0 9). 1. SEED SHAPE: 2	P. O. Box 959 Minneapolis, MN 55440	Code)	PVPO NUMBER	
1 Spherical (L/M, L/T, and T/M ratios = < 1.2) 2 Spherical Flattened (L/M ratio > 1.2: L/T ratio = < 1.2) 3 = Elongate (L/T ratio > 1.2: T/M = < 1.2) 4 = Elongate Flattened (L/T ratio > 1.2: T/M > 1.2) 4 = Elongate Flattened (L/T ratio > 1.2: T/M > 1.2) 2 SBED COAT COLOR: (Mature Seed) 1 1 Yellow 2 = Green 3 = Brown 4 = Black 5 = Other (Specify) 3 SEED COAT LUSTER: (Mature Hand Shelled Seed) 1 1 Dull (*Corsoy 79*; *Braxton*) 2 = Shiny (*Nebsoy*; *Gasoy 17*) 4 SEED SIZE: (Mature Seed) 1 4 Grams per 100 seeds 5 HILUM COLOR: (Mature Seed) 6 1 Buff 2 Yellow 3 = Brown 4 = Gray 5 = Imperfect Black 5 = Black 7 = Other (Specify) May contain up to 2% other hilum color. 6 COTYLEDON COLOR: (Mature Seed) 1 1 Yellow 2 = Green 7 SEED PROTEIN PEROXIDASE ACTIVITY: 2 1 Low 2 = High 2 = Type B (SP1 ^b) 3 Light Purple below cotyledons (*Beson'; *Pickett 71*) 4 Dark Purple extending to unifoliate leaves (*Hodgson*; *Coker Hampton 266A*) 10 LEAFLET SHAPE:	Choose the appropriate response which characterizes the		below. When the num	iber of significant digits
1 1 = Yellow 2 = Green 3 = Brown 4 = Black 5 = Other (Specify) 3. SEED COAT LUSTER: (Mature Hand Shelled Seed) 1 1 = Dull ('Corsov 79'; 'Braxton') 2 = Shiny ('Nebsoy'; 'Gasoy 17') 4. SEED SIZE: (Mature Seed) 1 4 Grams per 100 seeds 5. HILUM COLOR: (Mature Seed) 6 1 = Buff 2 = Yellow 3 = Brown 4 = Gray 5 = Imperfect Black 6 = Black 7 = Other (Specify) May contain up to 2% other hilum. Color. 6. COTYLEDON COLOR: (Mature Seed) 1 1 = Yellow 2 = Green 7. SEED PROTEIN PEROXIDASE ACTIVITY: 2 1 = Low 2 = High 8. SEED PROTEIN ELECTROPHORETIC BAND: 2 1 = Type A (Sp1ª) 2 = Type B (Sp1b) 9. HYPOCOTYL COLOR: 2 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A')	1 = Spherical (L/W, L/T, and T/W ratios = < 1.2)	W T T 2 = Spherical Flattened		_
3. SEED COAT LUSTER: (Mature Hand Shelled Seed) 1 1 = Dull ('Corsoy 79'; 'Braxton') 2 = Shiny ('Nebsoy'; 'Gasoy 17') 4. SEED SIZE: (Mature Seed) 1 4 Grams per 100 seeds 5. HILUM COLOR: (Mature Seed) 6 1 = Buff 2 = Yellow 3 = Brown 4 = Gray 5 = Imperfect Black 6 = Black 7 = Other (Specify) May contain up to 2% other hilum.color. 6. COTYLEDON COLOR: (Mature Seed) 1 1 = Yellow 2 = Green 7. SEED PROTEIN PEROXIDASE ACTIVITY: 2 1 = Low 2 = High 8. SEED PROTEIN ELECTROPHORETIC BAND: 2 1 = Type A (SP1*) 2 = Type B (SP1*) 9. HYPOCOTYL COLOR: 2 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Besson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A')		4- Pi-d	(0:6-1	
1 1 = Dull ('Corsov 79'; 'Braxton') 2 = Shiny ('Nebsoy'; 'Gasoy 17') 4. SEED SIZE: (Mature Seed) 5. HILUM COLOR: (Mature Seed) 6 1 = Buff 2 = Yellow 3 = Brown 4 = Gray 5 = Imperfect Black 6 = Black 7 = Other (Specify) May contain up to 2% other hilum.color. 6. COTYLEDON COLOR: (Mature Seed) 1 1 = Yellow 2 = Green 7. SEED PROTEIN PEROXIDASE ACTIVITY: 2 1 = Low 2 = High 8. SEED PROTEIN ELECTROPHORETIC BAND: 2 1 = Type A (SP18) 2 = Type B (SP18) 9. HYPOCOTYL COLOR: 2 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:	1 = Yellow 2 = Green 3 = Brown	4 = Black 5 = Other	(Specify)	
5. HILUM COLOR: (Mature Seed) 6. 1 = Buff		bsoy'; 'Gasoy 17')		
6 1 = Buff 2 = Yellow 3 = Brown 4 = Gray 5 = Imperfect Black 5 = Black 7 = Other (Specify) 6. COTYLEDON COLOR: (Mature Seed) 1 1 = Yellow 2 = Green 7. SEED PROTEIN PEROXIDASE ACTIVITY: 2 1 = Low 2 = High 8. SEED PROTEIN ELECTROPHORETIC BAND: 2 1 = Type A (SP1 ⁸) 2 = Type B (SP1 ^b) 9. HYPOCOTYL COLOR: 2 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A')				
May contain up to 2% other hilum.color. 6. COTYLEDON COLOR: (Mature Seed) 1	5. HILUM COLOR: (Mature Seed)			
1 = Yellow 2 = Green 7. SEED PROTEIN PEROXIDASE ACTIVITY: 2 1 = Low 2 = High 8. SEED PROTEIN ELECTROPHORETIC BAND: 2 1 = Type A (SP1 ⁸) 2 = Type B (SP1 ^b) 9. HYPOCOTYL COLOR: 2 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:			ack 6 = Black	7 = Other (Specify)
7. SEED PROTEIN PEROXIDASE ACTIVITY: 2 1 = Low 2 = High 8. SEED PROTEIN ELECTROPHORETIC BAND: 2 1 = Type A (SP1 ⁸) 2 = Type B (SP1 ^b) 9. HYPOCOTYL COLOR: 2 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:	6. COTYLEDON COLOR: (Mature Seed)			
8. SEED PROTEIN ELECTROPHORETIC BAND: 2 1 = Type A (SP1 ^a) 2 = Type B (SP1 ^b) 9. HYPOCOTYL COLOR: 2 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:	1 = Yellow 2 = Green		44	
8. SEED PROTEIN ELECTROPHORETIC BAND: 2 = Type B (SP1 ^b) 9. HYPOCOTYL COLOR: 1 = Green only ('Evans'; 'Davis')	7. SEED PROTEIN PEROXIDASE ACTIVITY:			
2 = Type B (SP1 ^b) 9. HYPOCOTYL COLOR: 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:	2 1 = Low 2 = High			
9. HYPOCOTYL COLOR: 1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:	8. SEED PROTEIN ELECTROPHORETIC BAND:			
1 = Green only ('Evans'; 'Davis') 2 = Green with bronze band below cotyledons ('Woodworth'; 'Tracy') 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:	2 = Type B (SP1 ^b)	,		
3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; 'Coker Hampton 266A') 10. LEAFLET SHAPE:	9. HYPOCOTYL COLOR:			
-	3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71	17)	'Woodworth'; 'Tracy')	
	10. LEAFLET SHAPE:	4-04-5		

		 -		, , , , , , ,	0.0
11. LEAF	LET SIZE:	.:			
2	1 = Small ('Amsoy 71'; 'A5312') 3 = Large ('Crawford'; 'Tracy')	2 = Medium ('Corso	y 79'; 'Gasoy 17')		
12. LEAF	COLOR:				•
2	1 ~ Light Green ('Weber'; 'York') 3 ~ Dark Green ('Gnome'; 'Tracy')	2 = Medium Green	('Corsoy 79'; 'Braxton')	· · · · · · · · · · · · · · · · · · ·	
13. FLOV	ER COLOR:				
1	1 = White 2 = Purple	3 = White with purple t	throat	<u></u>	
14, POD (COLOR:				
1	1 = Tan 2 = Brown	3 = Black	·		
15. PLAN	T PUBESCENCE COLOR:				
2	1 = Gray 2 = Brown (Tawny)				
16. PLAN	T TYPES:				
2	1 = Slender ('Essex'; 'Amsoy 71') 3 = Bushy ('Gnome'; 'Govan')	2 = Intermediate ('A	Amcor'; 'Braxton')		
17. PLAN	T HABIT:				· .
3	1 = Determinate ('Gnome'; 'Braxton') 3 = Indeterminate ('Nebsoy'; 'Improved P		te ('Will')		
TO MATE	RITY GROUP:				
5	1 = 000	4 = I 5 = II III 12 = IX 13 = X	· ·	IV 8 = V	
19. DISEA	SE REACTION: (Enter 0 = Not Tested; 1 =	Susceptible; 2 = Resistant)			
BACT	ERIAL DISEASES:			TIELL.	
	Bacterial Pustule (Xanthomonas phaseoli	var. sojensis)		RECEIV	50 K12
1	Bacterial Blight (Pseudomonas glycinea)			USDA A	MS E
一	Wildfire (Pseudomonas tabaci)	•		MAY 4 -	1990 D
FUNG/	AL DISEASES:	•		A Plant Vario	ary F2
	Brown Spot (Septoria glycines)			Prefection (Ofc S
	Frogeye Leaf Spot (Cercospora sojina)			TO THE	1110
	Race 1 Race 2 F	Race 4	Race 5	Other (Specify)	
	Target Spot (Corynespora cassiicola)				
	Downy Mildew (Peronospora trifoliorum y	/ar. manshurica)			
	Powdery Mildew (Microsphaera diffusa)				
- 2	Brown Stem Rot (Cephalosporium gregatu	m) (Horizontal Re	sistance based on	field observ	ation)
	Stem Canker (Diaporthe phaseolorum var.	caulivora)	Table 1		

19. DISEA	SE REACTION	: (Enter 0 = Not To	ested; 1 = Susceptib	le; 2 = Resistant) (C	ontinued)			
FUN	GAL DISEASES	S: (Continued)	_					
- 1	Pod and Stem	Blight (Diaporthe)	phaseolorum var; so	– ojae)				
1	Purple Seed S	tain <i>(Cercospora ki</i>	kuchii)					
	Rhizoctonia F	Root Rot <i>(Rhizocto</i>	nia solani) -					
	Phytophthora	Rot (Phytophthor	a megasperma var. se	ojae)				
1	Race 1	1 Race 2	1 Race 3	1 Race 4	1 Race 5	1 Race 6	1 Race	7
1	Race 8	1 Race 9	Other (Spec	cify)			· · · · · · · · · · · · · · · · · · ·	
VIR	AL DISEASES:							
	Bud Blight (To	obacco Ringspot V	irus)	,				
	Yellow Mosaid	: (Bean-Yellow Mos	aic Virus)					
	Cowpea Mosai	ic (Cowpea Chlorot	ic Virus)					
	Pod Mottle (B	ean Pod Mottle Vir	us}				e.	
	Seed Mottle (S	Soybean Mosaic Vir	us)					
NEM	IATODE DISEA	SES:	•					
18 .	Soybean Cyst	Nematode (Hetero	dera glycines)		·			
1	Race 1	1 Race 2	1 Race 3	1 Race 4	Other (S	pecify)		
	Lance Nemato	 ode (<i>Hopiolaimus C</i>	olombus)					
H		-	Meloidogyne incogn	nita)				
H			Meloidogyne Hapla)					
	Peanut Root K	Cnot Nematode (Me	loidogyne arenaria))				
	Reniform Nem	natode (<i>Rotylenchu</i>	lus reniformis)					
	OTHER DISE	ASE NOT ON FOR	M (Specify):				· · · · · · · · · · · · · · · · · · ·	·
		e de la compania de l La compania de la compania del compania de la compania de la compania del compania de la			•			
20. PHYSIC	DLOGICAL RES	SPONSES: (Enter (= Not Tested; 1 =	Susceptible; 2 = Resi	istant)			
1	Iron Chlorosis	on Calcareous Soil	Moderately	susceptible	or interm	ediate		
	Other (Specify) <u> </u>					· .	
21. INSEC	REACTION: (Enter 0 = Not Test	ed; 1 = Susceptible;	; 2 = Resistant)				
	Mexican Bean	Beetle <i>(Epilachna v</i>	arivestis)					
	Potato Leaf Ho	opper (Empoasca fa	ibae)					
)					· 	
22 INDIC4				S THAT SUBMITTE	D.	· · · · · · · · · · · · · · · · · · ·		· · · · · · · · · · · · · · · · · · ·
	RACTER		OF VARIETY		RACTER	NAM	E OF VARIETY	
Plant Sh		Pres			oat Luster		-30	
Leaf Sha	ape	S20-	-26	Seed Siz	ze .	S34	-19	
Leaf Co	lor -	A312	27	Seed Sh	ape	s ² 27	-10	
Leaf Siz	e	Pres	ston	Seedling	Pigmentation	Wil	liams 82	
							1000	
FORM LM	3S-470-57 (2-82	r)					en jako en ega 1919aa. Nederlah	Page 3 of 4

9000165

23. GIVE DATA FOR SUBMITTED AND SIMILAR STANDARD VARIETY.	
The standard variety.	

VARIETY	NO. OF DAYS		CM PLANT	LEAFLET SIZE		SEED CONTENT		SEED SIZE	NO.
	MATURITY		HEIGHT	CM Width	CM Length	% Protein	% Oil	G/100 SEEDS	SEEDS/ POD
Submitted	126	2.1	84	5.3	9.6	39.7	20.6	13.6	2-3
Preston Name of Similar Variety	123	2.2	85	5.2	9.8	39.6	21.0	14.1	2-3

PUBLICATIONS USEFUL AS REFERENCE AIDS FOR COMPLETING THIS FORM:

- 1. Caldwell, B.E., ed. 1973. Soybeans: Improvement, Production, and Uses. Amer. Soc. Agron. Monograph No. 16.
- 2. Buttery, B.R. and R.I. Buzzell. 1968. Peroxidase activity in seeds of soybean varieties. Crop Sci., 8: 722-725.
- 3. Hymowitz, T. 1973. Electrophoretic analysis of SBTI-A₂ in the USDA soybean germplasm collection. Crop Sci., 13: 420-421.
- 4. Payne, R.C. and L.F. Morris. 1976. Differentiation of soybean cultivars by seedling pigmentation patterns. J. Seed Technol. 1: 1-19.




EXHIBIT D

Additional Description of the Variety

Soybean variety \$28-18 is a mid to late Maturity Group II cultivar maturing between Preston and Pella 86. It exhibits long hypocotyl extension when planted in sand at 12 cm depth at 25° C. It exhibits normal tolerance to metribuzin herbicide.

EXHIBIT E

Statement of the Basis of Applicant's Ownership

Soybean variety S28-18 was developed by the Northrup King Co. soybean breeding staff from germplasm sources cited in Exhibit A of this application. Northrup King Co. believes that the variety is novel as defined in the Plant Variety Protection Act and, therefore, that Northrup King Co. is the sole owner of the variety.